

# Low cerebral blood flow is a risk factor for severe intraventricular haemorrhage

Judith H Meek, Lidia Tyszczyk, Clare E Elwell, J S Wyatt

## Abstract

**Aims**—To investigate the relation between cerebral blood flow on the first day of postnatal life and the severity of any subsequent germinal matrix haemorrhage–intraventricular haemorrhage (GMH–IVH).

**Methods**—Cerebral blood flow was measured in 24 babies during the first 24 hours of life using near infrared spectroscopy. Repeated cerebral ultrasound examination was performed to define the maximum extent of GMH–IVH. Infants were classified as: normal scan, minor periventricular haemorrhage (haemorrhage that resolved), or severe GMH–IVH (haemorrhage distending the ventricles, that progressed to either post haemorrhagic dilatation or porencephalic cyst formation).

**Results**—Cerebral blood flow was significantly lower in the infants with GMH–IVH (median 7.0 ml/100 g/min) than those without haemorrhage (median 12.2 ml/100 g/min), despite no difference in carbon dioxide tension and a higher mean arterial blood pressure. On subgroup analysis, those infants with severe GMH–IVH had the lowest cerebral blood flow.

**Conclusion**—A low cerebral blood flow on the first day of life is associated with the subsequent development of severe intraventricular haemorrhage.

(Arch Dis Child Fetal Neonatal Ed 1999;81:F15–F18)

Keywords: intraventricular haemorrhage; cerebral blood flow; cerebral ischaemia; near infrared spectroscopy

Germinal matrix haemorrhage–intraventricular haemorrhage (GMH–IVH) in preterm neonates carries an increased risk of neurodevelopmental morbidity if it is associated with either post haemorrhagic ventricular dilatation (PHD) or haemorrhagic parenchymal infarction (HPI).<sup>1</sup> These lesions may follow early abnormalities of cerebral blood flow (CBF), and the risk of GMH–IVH is greatest during the first three days of life.<sup>2</sup> This study aimed to relate CBF measured during the first 24 hours of life to the incidence and severity of any subsequent GMH–IVH.

## Methods

We studied extremely preterm infants on the first day of postnatal life after admission to the University College London Hospitals Trust neonatal intensive care unit. Twenty four infants were studied. Ethical permission for the

study was obtained from the local ethics committee, and informed consent was obtained from one or both parents of each infant before the study. Infants who had been electively delivered on the basis of abnormal antenatal umbilical artery pulsations measured by Doppler ultrasonography were excluded, because their cerebral haemodynamics were likely to have been abnormal. Due to methodological limitations we were unable to study infants ventilated in air, which excluded those ventilated using the high frequency oscillator, or infants requiring 100% oxygen.

The median gestational age was 26 weeks (range 24 to 31 weeks), and the median birth weight was 873 g (range 620 to 1470 g). All infants were appropriately grown for gestational age. All were ventilated with supplementary oxygen using intermittent positive pressure ventilation. Two were receiving infusions of dopamine and/or dobutamine to treat hypotension. Clinical details are given in table 1.

Cerebral blood flow was measured using near infrared spectroscopy (NIRS)<sup>3</sup> during the first day after delivery, at a median (range) age of 19 (5–24) hours. Measurements were obtained using a commercial spectrophotometer (NIRO500, Hamamatsu Photonics KK, Japan). Near infrared light from four laser diodes was transmitted via a fibre optic bundle to an “optode” on the temporo-parietal or frontal region of the infant’s head. A detecting optode, placed 4–5 cm away on the opposite side of the head, was connected to a second fibre optic bundle which conveyed the emerging light to the spectrophotometer. A computer converted the changes in optical density into changes in cerebral concentration of oxyhaemoglobin (HbO<sub>2</sub>), and deoxyhaemoglobin from an arbitrary baseline. Arterial oxygen saturation (SaO<sub>2</sub>), mean arterial pressure (MABP), partial pressure of carbon dioxide (PaCO<sub>2</sub>) and heart rate were simultaneously recorded.

Cerebral blood flow was determined from the Fick principle, using HbO<sub>2</sub> as an intravascular tracer. A rapid rise in SaO<sub>2</sub> was induced by a step increase in FIO<sub>2</sub>. Cerebral blood flow (ml/100 g/min) was then calculated from the integral of the saturation rise and the rate of rise of tracer concentration using the equation:

$$CBF = k \cdot \Delta [HbO_2] / (H \cdot \int \Delta SaO_2 \cdot dt)$$

where k is a constant (0.0614) incorporating brain tissue density and the molecular weight of haemoglobin and H is the large vessel haemoglobin concentration (g/dl).

Department of  
Paediatrics  
University College  
London Medical  
School  
WC1E 6JJ  
J H Meek  
L Tyszczyk  
J S Wyatt

Department of  
Medical Physics and  
Bioengineering  
C E Elwell

Correspondence to:  
Dr J H Meek.  
Email: [j.meek@ucl.ac.uk](mailto:j.meek@ucl.ac.uk)

Accepted 9 December 1998

Table 1 Clinical details and ultrasound diagnoses of infants studied

Group	Infant	Gestational age	Birth weight SDS	Diagnosis	Inotropes	Clinical outcome	Cranial ultrasound diagnosis, with day haemorrhage first noted
1	1	24	1.29	Mild HMD	No	No major problems	Normal
	2	28	0.50	PROM, mild HMD	No	No major problems	Normal
	3	25	-0.07	HMD	No	No major problems	Normal
	4	29	-1.34	Triplet I	No	No major problems	Normal
	5	27	-0.42	Twin I	Dopamine at 10 µg/kg/min	No major problems	Normal
	6	27	-0.38	HMD	No	No major problems	Normal
	7	24	-1.07	HMD	No	No major problems	Normal
	8	28	-0.84	Severe HMD and PIE	No	Mild spastic diplegia	Normal
	9	24	-0.48	Severe HMD and PIE	No	Died day 5, PIE	Normal
	10	25	-1.07	HMD	No	No major problems	Normal
	11	26	1.29	Severe HMD	No	No major problems	Normal
	12	28	-0.97	Mild HMD	No	No major problems	Normal
	13	28	-0.91	Mild HMD	No	No major problems	Normal
	14	31	-0.13	Moderate HMD	No	Emigrated, no follow-up	Normal
	15	30	0.57	Mild HMD	No	No major problems	Normal
	16	27	-0.94	Maternal PIH, mild HMD	No	No major problems	Normal
	17	25	1.33	Maternal streptococcal chorioamnionitis	No	No major problems	No haemorrhage, late small periventricular cyst
2	18	25	0.35	Maternal candidiasis, mild HMD	No	No major problems	Left GLH, right small IVH day 1
3	19	26	-0.58	Maternal PIH, mild HMD	No	No major problems	Right GLH day 2
	20	29	0.68	Twin 2, mild HMD	No	Diplegia, esotropia	Bilateral large IVH with PHD day 2
3	21	26	-0.44	Severe HMD and PIE	No	Hemiplegia, developmental delay	Left HPI, leading to porencephalic cyst
	21	26	-0.44	Severe HMD and PIE	No	developmental delay	Right small IVH day 2
3	22	27	0.72	HMD	No	Convulsions	Left HPI, leading to porencephalic cyst
	22	27	0.72	HMD	No	Hemiplegia, asthma	Right large IVH day 1
3	23	24	-0.14	Twin 1, HMD	No	Developmental delay	Left HPI and right large IVH, PHD day 2
	24	24	0.04	Twin 2, severe HMD, renal failure	Dopamine, dobutamine	Died day 5	Left large IVH, right HPI, PHD day 3

Abbreviations: HMD, hyaline membrane disease; PIE, pulmonary interstitial emphysema.

For each infant six measurements of CBF were attempted. Only those measurements were accepted that met previously established criteria for the stability of parameters such as the PaCO<sub>2</sub>, MABP, and [Hb + HbO<sub>2</sub>] during the measurement, and the stability of the SaO<sub>2</sub> baseline before the saturation swing.<sup>4</sup> The mean value of the measurements which met the acceptance criteria was used for analysis. Between two and six successful measurements were achieved for each infant.

Cerebral ultrasound examination was performed daily for the first week of life, and then at least once a week until term or death. Scans were recorded and carefully classified by independent observers who were blind to the cerebral blood flow results. All infants had a normal appearance on cranial ultrasound examination at the time of study.

Infants were ascribed to outcome groups according to the maximum extent of the appearance of their cranial ultrasound scans. Group 1 had normal scans and group 2 had evidence of minor GMH-IVH which later resolved leaving a normal scan appearance. Group 3 had large GMH-IVH, distending the ventricles, progressing either to PHD or HPI with subsequent porencephalic cyst formation. The CBF, PaCO<sub>2</sub>, MABP and age of each group at the time of study were compared with the others using the Kruskal-Wallis one way analysis of variance (SPSS). As the numbers of infants in groups 2 and 3 were small, the same parameters were compared between the normal group (1) and all infants with GMH-IVH (groups 2 and 3 combined) using the Mann-Whitney test.

## Results

Seventeen of the 24 infants had normal appearances on cranial ultrasound examination (group 1). Two had minor haemorrhages (group 2),

and one of these developed a small unilateral parenchymal cyst. Five had severe haemorrhages (group 3). The individual diagnoses and times of onset of the haemorrhage for each infant in groups 2 and 3 are shown in table 1. The timings of the maximum extent of the lesions ranged between 2 and 14 days. The median (range) CBF of infants in group 1 was 12.2 ml/100 g/min (6.1–24.7), in group 2, 12.0 ml/100 g/min (10.3–13.8), and in group 3 was 5.8 ml/100 g/min (3.2–7.5). Cerebral blood flow was significantly lower in infants with any GMH-IVH (groups 2 and 3) than in infants without lesions (group 1) ( $p < 0.04$ , Mann-Whitney U test), and was significantly lower in group 3 on subgroup analysis ( $p < 0.01$ , Mann-Whitney U test), but not in group 2. There was a significant decrease in CBF with group number ( $p=0.02$  by Kruskal-Wallis one way Anova). Figure 1 shows the distribution of CBF over the three groups. There was no significant difference between the PaCO<sub>2</sub>, gestational age, age at the time of study, inspired oxygen fraction or mean airway pressure of each group. There was a positive trend between MABP and group number ( $p = 0.05$ , Kruskal-Wallis), and MABP values in group 3 were significantly higher than those in group 1 ( $p < 0.02$  Mann-Whitney U test). These results are summarised in table 2. Figure 2 shows the distribution of MABP over the three groups.

## Discussion

Cerebral blood flow measured during the first 24 hours of life was lower in infants who subsequently developed severe GMH-IVH compared with those who did not develop haemorrhages.

All clinically feasible methods for CBF measurement in newborn infants have methodological limitations. The HbO<sub>2</sub> technique for




Figure 1 Comparison of cerebral blood flow for the three groups.




Figure 2 Comparison of MABP for the three groups.

measuring CBF has been validated against the intravenous  $^{133}\text{Xe}$  clearance method.<sup>5,6</sup> The assumptions underlying the method have been discussed elsewhere.<sup>7</sup> It is reasonable to assume that our results reflect genuine differences in cerebral perfusion between the different study groups.

The range of CBF values in each group was wide. This may, in part, reflect the fact that measurements were performed over short time periods, and were obtained at different times within the first day of life. Repeated measurements of CBF standardised for time over the first 24 hours would probably yield a more detailed picture of changes in cerebral perfusion which precede the development of GMH-IVH. There was no significant difference in  $\text{PaCO}_2$ , gestational age, or postnatal age at the time of study between the groups. MABP was significantly higher in the infants who developed severe GMH-IVH compared with those without haemorrhage.

Table 2 Results for each group (median and range)

Group	CBF (ml/100 g/min)	MABP (mm Hg)	$\text{PaCO}_2$ (kPa)	Age at study (hours)	Gestational age (weeks)
1	12.1 (6.1–24.7)	29.5 (25–40)	5.6 (3.8–7.6)	15 (6–24)	27 (24–31)
2	12.0 (10.3–13.8)	31.0 (27–35)	5.2 (5.2–6.3)	12.5 (12–13)	25.5 (25–26)
3	5.8 (3.2–7.5)	35.0 (35–43)	5.6 (4.9–5.8)	20 (8–17)	26 (24–29)

Since the seminal contribution of Pape and Wigglesworth,<sup>8</sup> cerebral ischaemia has been suggested as an important factor in the pathophysiology of GMH-IVH. Hypoperfusion of the germinal matrix is thought to lead to an impairment in the integrity of the vascular endothelium. A subsequent period of reperfusion leads to an abrupt increase in intravascular pressure, causing haemorrhage within the germinal matrix. It is thought that the extent and position of the haemorrhage determine the likelihood of major complications such as progression to HPI or PHD. The vulnerability of the germinal matrix to hypoperfusion may be due to its watershed vascular supply, or to the higher metabolic rate of its endothelium.<sup>9</sup> Low CBF during the first day of life may damage the germinal matrix in a way that preconditions it to haemorrhage.

The concept that GMH-IVH was due to an ischaemia/reperfusion sequence was reinforced by the work of Goddard-Finegold and colleagues in a newborn beagle model. They showed that intraventricular haemorrhage occurred in animals exposed to rapid volume expansion, only if it was preceded by hypotension.<sup>10</sup>

GMH-IVH is associated with several factors, especially prematurity, the presence of lung disease and a requirement for mechanical ventilation.<sup>11,12</sup> The association may be mediated by fluctuations in arterial oxygen and carbon dioxide tensions, and in arterial pressure, leading to periods of intermittent ischaemia interspersed with episodes where reperfusion occurs. This may be exacerbated by immaturity of cerebral autoregulation.<sup>13,14</sup> Mechanical ventilation and pneumothorax are also associated with fluctuations in intrathoracic pressure which may be transmitted to the cerebral venous system.

Perlman and colleagues, using Doppler ultrasound, showed an association between fluctuations in blood flow velocity in the anterior cerebral artery and the subsequent development of intraventricular haemorrhage,<sup>15</sup> but other workers have failed to reproduce their observations.<sup>16</sup>

Although rapid fluctuations in cerebral perfusion may predispose to haemorrhage, it is also likely that a prolonged episodes of hypoperfusion may cause vascular injury and programme the germinal matrix towards subsequent haemorrhage.

Ment *et al* have shown that infants who sustain GMH-IVH during the first 6 hours of life have lower CBF (measured by inhalation  $^{133}\text{Xe}$  clearance) than both those who subsequently develop haemorrhage, and those with normal scans.<sup>17</sup> They have suggested that the lower CBF may be due to the presence of the haemorrhage itself.

The possibility that a cerebral “steal” phenomenon through a patent ductus arteriosus may contribute to the pathogenesis of GMH-IVH has been proposed by Evans and Kluckow,<sup>18,19</sup> based on observations that large early PDA shunts are significantly associated with GMH-IVH,<sup>20</sup> and that a PDA can be

associated with reversed flow velocity in the anterior cerebral arteries,<sup>21</sup> and the descending aorta.<sup>22</sup> They measured superior vena cava (SVC) flow<sup>19</sup> as well as the degree of ductal and atrial shunting using Doppler echocardiography at 5, 12, 24 and 48 hours of age in a group of very preterm babies. They found that lower SVC flow at 5 and 12 hours of age was significantly related to subsequent haemorrhage, and was associated with large ductal diameter. SVC flow tended to increase after 12 hours of age coinciding with the appearance of cranial ultrasound evidence of haemorrhage.

The concept that a cerebral steal phenomenon precedes the development of GMH-IVH may explain the reduction in the incidence and severity of haemorrhage following the early administration of indomethacin.<sup>23</sup> This effect contrasts with the decrease in CBF which is seen when indomethacin is used for late closure of PDAs.<sup>24</sup> By combining measurements of CBF and ductal shunting during the first few hours of life, it may be possible to identify the subgroup of infants who would benefit from treatment with indomethacin, and avoid subjecting the remaining infants to its potential adverse effects.

Further evidence suggests that a PDA, rather than being a cause of cerebral ischaemia, may be a compounding influence. Evans and Kluckow<sup>25</sup> have shown an inverse relation between SVC flow and upper body vascular resistance. Low CBF resulting from the immature myocardium adapting to extrauterine vascular resistances, is further compromised by ductal steal.

Our findings are consistent with the hypothesis that cerebral ischaemia is an important predisposing factor in the development of severe GMH-IVH. Interestingly, there was no difference in PaCO<sub>2</sub> among the study groups, and MABP was higher in those infants with low CBF. It is not clear, therefore, why CBF was impaired in the adverse outcome group. Although we did not perform echocardiography at the time of study, and no infants had obvious clinical evidence of ductal shunting, a cerebral steal phenomenon might have contributed to the low CBF. Further studies are urgently required to investigate the relation between clinically silent ductal shunting<sup>26</sup> and cerebral perfusion in the critical period immediately after birth when programming of GMH-IVH may occur.

This study was funded by the Medical Research Council. We thank Dr Ann Stewart, Professor David Edwards, Dr David McCormick, Dr Vincent Kirkbride and Ms Jan Townsend for their contribution to this work. We are grateful to Dr Jane Hawdon and Dr Nick Evans for valuable discussions.

- Roth SC, Baudin J, McCormick DC, *et al*. Relation between ultrasound appearance of the brain of very preterm infants and neurodevelopmental impairment at eight years. *Dev Med Child Neurol* 1993;35:755-68.
- Thorburn RJ, Lipscomb AP, Stewart AL, Reynolds EOR, Hope PL. Timing and antecedents of periventricular haemorrhage and of cerebral atrophy in very preterm infants. *Early Hum Dev* 1982;7:221-38.
- Edwards AD, Wyatt JS, Richardson C, Delpy DT, Cope M, Reynolds EOR. Cotside measurement of cerebral blood flow in ill newborn infants by near infrared spectroscopy. *Lancet* 1988;ii: 770-1.
- Elwell CE. *A practical user's guide to near infrared spectroscopy*. Japan: Hamamatsu Photonics, 1995.
- Bucher HU, Lipp AF, Duc D, Edwards AD. Comparison between <sup>135</sup>Xenon clearance and near infrared spectroscopy for estimation of cerebral blood flow in sick preterm infants. *Pediatr Res* 1993; 33:56-60.
- Skov L, Pryds O, Greisen G. Estimating cerebral blood flow in newborn infants: comparison of near infrared spectroscopy and <sup>133</sup>Xe clearance. *Pediatr Res* 1991;30:570-3.
- Elwell CE, Cope M, Edwards AD, Wyatt JS, Reynolds EOR, Delpy DT. Measurement of cerebral blood flow in adult humans using near infrared spectroscopy - methodology and possible errors. *Adv Exp Med Biol* 1992;317:235-45.
- Pape KE, Wigglesworth JS. Haemorrhage, ischaemia and the perinatal brain. In: *Clinics in Developmental Medicine* 69/70 London: SIMP/Heinemann, 1979.
- Volpe JJ. *Neurology of the Newborn*. 3rd Edn. Philadelphia: W B Saunders & Co, 1995: 418.
- Goddard-Finegold J, Armstrong D, Zeller RS. Intraventricular haemorrhage following volume expansion after hypovolaemic hypotension in the newborn beagle. *J Pediatr* 1982;100:796-9.
- Levene MI, Fawer CL, Lamont RF. Risk factors in the development of intraventricular haemorrhage in the preterm neonate. *Arch Dis Child* 1982; 57:410-17.
- Harrison VC, Heese H, Klein M. Intracranial haemorrhage associated with hyaline membrane disease. *Arch Dis Child* 1968;43:116-20.
- Pryds O. Control of the cerebral circulation in the high-risk neonate. *Ann Neurol* 1991 30:321-9.
- Miall-Allen VM, De Vries LS, Whitelaw AGL. Mean arterial blood pressure and neonatal cerebral lesions. *Arch Dis Child* 1987;62:1068-9.
- Perlman JM, McMenamin JB, Volpe JJ. Fluctuating cerebral blood-flow velocity in respiratory distress syndrome. Relation to the development of intraventricular haemorrhage. *N Engl J Med* 1983;309:204-9.
- Miall-Allen VM, de Vries LS, Dubowitz LMS, Whitelaw AGL. Blood pressure fluctuations and intraventricular haemorrhage in the preterm infant of less than 31 weeks' gestation. *Pediatrics* 1989;83:657-61.
- Ment LR, Duncan CC, Ehrenkranz RA, *et al*. Intraventricular haemorrhage in the preterm neonate: timing and cerebral blood flow changes. *J Pediatr* 1984;104:419-25.
- Evans NJ, Kluckow M. The haemodynamic basis of intraventricular haemorrhage (IVH). *Pediatr Res* 1997;43:173A.
- Evans NJ, Kluckow M. Early determination of right and left ventricular output in ventilated preterm infants. *Arch Dis Child* 1996;74:F88-94.
- Evans N, Kluckow M. Early ductal shunting and intraventricular haemorrhage in ventilated preterm infants. *Arch Dis Child* 1996; 75:F183-6.
- Perlman JM, Hill A, Volpe JJ. The effect of patent ductus arteriosus on flow velocity in the anterior cerebral arteries ductal steal in the premature infant. *J Pediatr* 1981; 99:767-71.
- Lipman A, Serwer GA, Brazy JE. Abnormal cerebral haemodynamics in preterm infants with patent ductus arteriosus. *Pediatr Res* 1982; 69:778-81.
- Ment LR, Oh W, Ehrenkranz RA, *et al*. Low dose indomethacin and prevention of intraventricular haemorrhage: a multicenter randomized trial. *Pediatrics* 1994;93:543-50.
- Edwards AD, Wyatt JS, Richardson C, *et al*. Effects of indomethacin on cerebral haemodynamics in very preterm infants. *Lancet* 1990;335:1491-5.
- Evans N, Kluckow M. Low upper body and cerebral blood flow in the preterm infant. *Pediatr Res* 1998;43:173A.
- Skelton R, Evans N, Smythe J. A blinded comparison of clinical and echocardiographic evaluation of the preterm infant for patent ductus arteriosus. *J Paediatr Child Health* 1994;30:406-11.



## Low cerebral blood flow is a risk factor for severe intraventricular haemorrhage

Judith H Meek, Lidia Tyszczuk, Clare E Elwell, et al.

*Arch Dis Child Fetal Neonatal Ed* 1999 81: F15-F18  
doi: 10.1136/fn.81.1.F15

---

Updated information and services can be found at:  
<http://fn.bmj.com/content/81/1/F15.full.html>

---

### References

*These include:*

This article cites 23 articles, 6 of which can be accessed free at:  
<http://fn.bmj.com/content/81/1/F15.full.html#ref-list-1>

Article cited in:  
<http://fn.bmj.com/content/81/1/F15.full.html#related-urls>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

[Hypertension](#) (173 articles)  
[Clinical diagnostic tests](#) (437 articles)  
[Oncology](#) (127 articles)  
[Radiology](#) (420 articles)  
[Radiology \(diagnostics\)](#) (389 articles)

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>