

CURRENT TOPIC

Low birthweight and adult insulin resistance: the “catch-up growth” hypothesis

Stefano Cianfarani, Daniela Germani, Francesco Branca

Fetal origin of adult diseases

Epidemiological studies have shown a close correlation between intrauterine growth retardation (IUGR) and the onset of insulin resistance, non-insulin-dependent diabetes mellitus (NIDDM), hypertension, hyperlipidaemia, and cardiovascular diseases in adult life.¹⁻⁶ To explain this association, the concept of re-programming has been introduced: intrauterine exposure to insufficient nutrient supply during critical periods of fetal life would permanently affect the development and function of the endocrine system, leading to metabolic changes, including reduced insulin sensitivity.^{5,6}

Although knowledge of the mechanisms involved in the re-programming process might allow new strategies for early prevention of long term metabolic disturbances to be developed, the pathophysiological link between fetal growth impairment and adult diseases is still unclear.

The “thrifty phenotype” and the “fetal salvage” hypotheses

In 1992 Hales and Barker⁷ proposed the model of the “thrifty phenotype,” suggesting that intrauterine malnutrition would lead to insulin resistance and decreased β cell mass, thus predisposing to NIDDM. According to this hypothesis, the endocrine alterations induced by intrauterine malnutrition are intended to divert the limited nutrient supply to maintain survival and development of vital organs, such as the brain, at the expense of growth.

More recently, the “fetal salvage” hypothesis has been formulated.⁸ The finding that prepubertal IUGR children show a far greater insulin response than normal birthweight children, challenges the previously proposed β cell hypoplasia.⁹ The “fetal salvage” model suggests that the malnourished fetus develops peripheral insulin resistance which allows a redistribution of nutrients, such as glucose, in favour of essential organs. This then leads to a permanent reduction in skeletal muscle glucose transporter number or function. This reduced peripheral insulin sensitivity stimulates β cells to produce larger amounts of insulin to achieve normal glycaemia and would lead to eventual β cell exhaustion. This hypothesis is supported by studies in animal models showing reduced glucose transporter protein concentrations in

skeletal muscles of IUGR fetuses and normal concentrations in the brain.^{10,11}

Insulin-like growth factors: more than modulators of growth

The insulin-like growth factors (IGFs)-I and -II are structurally related to insulin, bind to specific receptors (type 1 and 2 IGF receptors) and, with lower affinity, to insulin receptors. IGFs circulate in blood bound to a family of at least 10 different binding proteins (IGFBPs) that regulate the biological activities of the IGFs, also exerting IGF independent actions.¹²⁻¹⁴ The affinity of IGFBPs for IGFs can be reduced by specific IGFBP proteases which, by fragmentation of IGFBPs, lower the binding of IGFs and increase IGF bioavailability.¹⁵ IGFs exert primarily growth promoting and insulin-like actions. IGF-I administration to humans affects glucose metabolism, stimulating glucose uptake and leading to hypoglycaemia. At higher concentrations, IGFs can bind to insulin receptors and the administration of IGFBP-1, a known inhibitor of IGF biological actions, increases glucose concentrations.¹⁶

A role for IGFs in fetal growth is suggested by numerous observations. A close correlation between umbilical cord serum IGF-I concentrations and birthweight has repeatedly been reported.¹⁷⁻¹⁹ IGF-I and IGFBP-3 concentrations are reduced in IUGR children, but IGFBP-1 and IGFBP-2 concentrations are increased in samples obtained in utero and cord serum.^{20,21} Targeted mutagenesis of the genes encoding IGF-I and -II and type 1 IGF receptor induces profound embryonic, fetal, and postnatal growth retardation in mice.^{22,23} The first patient with severe IUGR and postnatal growth retardation caused by a deletion of IGF-I gene has been described.²⁴ Finally, the increased IGFBP-3 proteolytic activity found in mothers with multiple fetuses or fetuses affected by uteroplacental insufficiency suggests a plausible mechanism for the increased IGF bioavailability whenever fetal growth is threatened.²⁵

We recently observed the presence of at least two different IGFBP-3 proteases in the serum of IUGR children, the role of which might be the amplification of the IGF growth promoting action which determines early postnatal “catch-up” growth in most IUGR infants.²⁶

Laboratory of
Paediatric
Endocrinology
“Tor Vergata”
University
via di Tor Vergata 135
00133-Rome
Italy
S Cianfarani
D Germani

National Institute of
Nutrition
Rome, Italy
F Branca

Correspondence to:
Dr S Cianfarani.
Email:
stefano.cianfarani@uniroma2.it




Figure 1 Schematic representation of "catch-up growth" hypothesis. Adequate postnatal nutrient supply activates IGF system, leading to catch-up growth and insulin resistance. IGF: insulin-like growth factor; IGFBP-3: insulin-like growth factor binding protein-3.

The "catch-up growth" hypothesis

IUGR children show a rearrangement of the endocrine system at birth, having low concentrations of insulin, IGF-I, IGFBP-3, and high concentrations of growth hormone, IGFBP-1, and IGFBP-2. Normalisation of these parameters occurs during the first trimester of postnatal life.^{26, 27} It is conceivable that tissues chronically depleted of insulin and IGF-I during fetal life and suddenly exposed to increased concentrations of the two hormones shortly after birth, may counteract their additive insulin-like actions by developing insulin resistance as a metabolic defence mechanism to protect the organism from hypoglycaemia. According to this model, the crucial time for the development of long term consequences is early postnatal life when catch-up growth occurs in around 80% of IUGR children.²⁸ Consistent with our hypothesis, data from animals have shown that when fetal growth impairment is followed by catch-up growth postnatally, the lifespan is significantly shortened.²⁹ Moreover, the glucose induced insulin response in infants with catch-up growth is higher than that in children without significant catch-up growth.³⁰ Therefore, those IUGR infants who show early and complete recovery from intrauterine growth retardation would be at higher risk for the occurrence of metabolic disturbances like insulin resistance and NIDDM in adulthood (fig 1).

In addition, the recent demonstration of the ability of IGFBP-3 fragments generated by proteolytic cleavage to bind insulin, inhibiting binding and activation of insulin receptor,³¹ suggests that the increased IGFBP-3 proteolytic activity observed in the first months of life of IUGR children might contribute to the development of insulin resistance (fig 1).

Finally, Whitaker and colleagues have recently reported that an early "adiposity rebound," the point of maximal leanness usually achieved at 5 to 6 years of age, is associated with an increased risk of adult obesity.³² Early "adiposity rebound" has been suggested as constituting a marker for growth acceleration

and cell hyperplasia³³ which are, at least transiently, exaggerated during catch-up growth in IUGR infants who therefore might have an additional risk of developing insulin resistance in adulthood.

In conclusion, we speculate that the tremendous effort to recover height shortly after birth, involving overactivation of the IGF system kept quiescent during intrauterine life to divert the restricted nutrient supply from growth to survival of vital organs, induces a metabolic adaptation with long term effects. The secondary insulin resistance, when associated with other risk factors such as genetic predisposition or obesity, may eventually lead to NIDDM which would thus represent a potential long term consequence of catch-up growth in IUGR children.

- Barker DJP, Winter PD, Osmond C, Margetts B. Weight in infancy and death from ischaemic heart disease. *Lancet* 1989;2271:577-80.
- Law CM, Barker DJP, Bull AR, Osmond C. Maternal and fetal influences on blood pressure. *Arch Dis Child* 1991;66:1291-5.
- Robinson R, Walton R, Clark P, Barker DJP, Hales C, Osmond C. The relation of fetal growth to plasma glucose in young man. *Diabetologia* 1992;35:444-6.
- Barker DJP. Fetal origins of coronary heart disease. *BMJ* 1995;311:171-4.
- Barker DJP. Intrauterine programming of coronary heart disease and stroke. *Acta Paediatr* 1997;423(Suppl):178-82.
- Ravelli ACJ, van der Meulen JHP, Michels RPJ, et al. Glucose tolerance in adults after prenatal exposure to famine. *Lancet* 1998;351:173-7.
- Hales CN, Barker DJP. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 1992;35:595-601.
- Hofman PL, Cutfield WS, Robinson EM, et al. Insulin resistance in short children with intrauterine growth retardation. *J Clin Endocrinol Metab* 1997;82:402-6.
- Barker DJP, Hales CN, Fall CH, Osmond C, Phipps K, Clark PM. Type 2 (non-insulin-dependent) diabetes mellitus, hypertension, and hyperlipidaemia (syndrome X): relation to reduced fetal growth. *Diabetologia* 1993;36:62-7.
- Simmons RA, Gounis AS, Bangalora SA, Ogata ES. Intrauterine growth retardation: fetal glucose transport is diminished in lung but spared in brain. *Pediatr Res* 1985;31:59-63.
- Simmons RA, Flozak AS, Ogata ES. The effect of insulin, and insulin-like growth factor 1 on glucose transport in normal and small for gestational age fetal rats. *Endocrinology* 1993;133:1361-8.
- Cianfarani S, Holly JMP. Somatomedin-binding proteins: what role do they play in the growth process? *Eur J Pediatr* 1989;149:76-9.
- Jones JI, Clemmons DR. Insulin-like growth factors and their binding proteins: biological actions. *Endocrin Rev* 1995;16:3-34.
- Kim H, Nagalla SR, Oh Y, et al. Identification of a family of low-affinity insulin-like growth factor binding proteins (IGFBPs): characterization of connective tissue growth factor as a member of the IGFBP superfamily. *Proc Natl Acad Sci USA* 1997;94:12981-6.
- Rajaram S, Baylink DJ, Mohan S. Insulin-like growth factor-binding proteins in serum and other biological fluids: regulation and functions. *Endocrin Rev* 1997;18:801-31.
- Lewitt MS, Denyer GS, Cooney GJ, Baxter RC. Insulin-like growth factor-binding protein-1 modulates blood glucose levels. *Endocrinology* 1991;129:2254-6.
- Gluckman PD, Johnson-Barrett JJ, Butler JH, Edgar BW, Gunn TR. Studies of insulin-like growth factor-I and -II by specific radioligand assays in umbilical cord blood. *Clin Endocrinol (Oxf.)* 1983;19:405-13.
- Lassarre C, Hardouin S, Daffos F, Forestier F, Frankenne F, Binoux M. Serum insulin-like growth factors and insulin-like growth factor binding proteins in the human fetus. Relationships with growth in normal subjects and in subjects with intrauterine growth retardation. *Pediatr Res* 1991;29:219-25.
- Verhaeghe J, Van Bree R, Van Herck E, Laureys J, Bouillon R, Van Assche A. C-peptide, insulin-like growth factors I and II and insulin-like growth factor binding protein-1 in umbilical cord serum: correlations with birth weight. *Am J Obstet Gynaecol* 1993;169:89-97.
- Giudice LC, de Zegher F, Gargosky SE, et al. Insulin-like growth factors and their binding proteins in the term and preterm human fetus and neonate with normal and extremes of intrauterine growth. *J Clin Endocrinol Metab* 1995;80:1548-55.

- 21 Hills FA, English J, Chard T. Circulating levels of IGF-I and IGF-binding protein-1 throughout pregnancy: relation to birth weight and maternal weight. *J Endocrinol* 1996;**148**:303–9.
- 22 Liu J-P, Baker J, Perkins AS, Robertson EJ, Efstratiadis A. Mice carrying null mutations of the genes encoding insulin-like growth factor-I (Igf-1) and type-1 IGF receptor (Igf1r). *Cell* 1993;**75**:59–72.
- 23 Baker J, Liu J-P, Robertson EJ, Efstratiadis A. Role of insulin-like growth factors in embryonic and postnatal growth. *Cell* 1993;**75**:73–82.
- 24 Woods KA, Camacho-Hubner C, Savage MO, Clark AJL. Intrauterine growth retardation and postnatal growth failure associated with deletion of the insulin-like growth factor I gene. *N Engl J Med* 1996;**335**:1363–7.
- 25 Langford KS, Nicolaidis KH, Jones J, Abbas A, McGregor AM, Miell JP. Serum insulin-like growth factor binding protein-3 (IGFBP-3) levels and IGFBP-3 protease activity in normal, abnormal, and multiple human pregnancy. *J Clin Endocrinol Metab* 1995;**80**:21–7.
- 26 Cianfarani S, Germani D, Rossi P, *et al*. Intrauterine growth retardation (IUGR): evidence for the activation of the IGF-related growth promoting machinery and the presence of a cation-independent IGFBP-3 proteolytic activity by two months of life. *Pediatr Res* 1998;**44**:374–80.
- 27 Leger J, Noel M, Limal JM, Czernichow P. Growth factors and intrauterine growth retardation. II. Serum growth hormone, insulin-like growth factor-I, and IGF-binding protein-3 levels in children with intrauterine growth retardation compared with normal control subjects: prospective study from birth to two years of age. *Pediatr Res* 1996;**40**:101–7.
- 28 Karlberg J, Albertsson-Wikland K. Growth in full-term small-for-gestational-age infants: from birth to final height. *Pediatr Res* 1995;**38**:733–9.
- 29 Hales CN, Desai M, Ozanne SE, Crowther NJ. Fishing in the stream of diabetes: from measuring insulin to the control of fetal organogenesis. *Biochem Soc Trans* 1996;**24**:341–50.
- 30 Colle E, Schiff D, Andrew G, Bauer CB, Fitzhardinge P. Insulin responses during catch-up growth of infants who were small for gestational age. *Pediatrics* 1976;**57**:363–71.
- 31 Yamanaka Y, Wilson EM, Rosenfeld RG, Oh Y. Inhibition of insulin receptor activation by insulin-like growth factor binding proteins. *J Biol Chem* 1997;**272**:30729–34.
- 32 Whitaker RC, Pepe MS, Wright JA, *et al*. Early adiposity rebound and the risk of adult obesity. *Pediatrics* 1998;**101**:E5.
- 33 Rolland-Cachera MF, Deheeger M, Bellisle F, *et al*. Adiposity rebound in children: a simple indicator for predicting obesity. *Am J Clin Nutr* 1984;**39**:129–35.



Low birthweight and adult insulin resistance: the "catch-up growth" hypothesis

Stefano Cianfarani, Daniela Germani and Francesco Branca

Arch Dis Child Fetal Neonatal Ed 1999 81: F71-F73

doi: 10.1136/fn.81.1.F71

Updated information and services can be found at:

<http://fn.bmj.com/content/81/1/F71.full.html>

References

These include:

This article cites 33 articles, 14 of which can be accessed free at:

<http://fn.bmj.com/content/81/1/F71.full.html#ref-list-1>

Article cited in:

<http://fn.bmj.com/content/81/1/F71.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

- [Drugs: CNS \(not psychiatric\)](#) (122 articles)
- [Drugs: endocrine system](#) (61 articles)
- [Diabetes](#) (88 articles)
- [Pregnancy](#) (844 articles)
- [Reproductive medicine](#) (795 articles)
- [Metabolic disorders](#) (186 articles)
- [Child health](#) (992 articles)
- [Lipid disorders](#) (6 articles)
- [Epidemiologic studies](#) (529 articles)
- [Hypertension](#) (173 articles)
- [Drugs: cardiovascular system](#) (172 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>